



# Search for Low-Mass WIMPs with SuperCDMS

Lauren Hsu

*for SuperCDMS*

*Fermilab Center for Particle Astrophysics*

**Joint Experimental-Theoretical Physics Seminar**

*Fermilab, March 14 2014*

# The dark matter problem

# An 80-year old puzzle



***In 1933:***


Fritz Zwicky analyzed velocity dispersion in Coma Cluster



Individual galaxies move too fast for a bound system...

***Posited existence of unseen matter in the cluster  
and named it “dark matter”***

# The Modern View of Dark Matter




## ***What we know:***

It's stable, cold, gravitationally interacting, non-baryonic, interacts little with itself (or not at all), composes  $\sim 80\%$  of matter in the Universe...

## ***But:***

If dark matter is composed of elementary particles, none in the Standard Model fits !

# WIMPs 101





"I THINK YOU SHOULD BE MORE EXPLICIT HERE IN STEP TWO."


# The Weakly Interacting Massive Particle

## *The WIMP "Miracle"*


Particles with mass and couplings at the weak scale yield cross sections that correspond to ~correct relic density of CDM



# How to detect WIMPs




*Relic  
annihilation in  
the cosmos*  
**INDIRECT  
DETECTION**



*man-made COLLIDER  
production*

*M. Attisha*




*Relic WIMP-  
nucleon elastic  
scattering*  
**DIRECT  
DETECTION**

# Consider the relic WIMP distribution

*Observed energy spectrum & rate depend on WIMP distribution in dark matter halo*

- Dark matter is distributed in a large extended, spherical halo around the Milky Way
- For comparison of direct detection experiments, assume an isothermal Maxwell-Boltzmann velocity distribution, with **width = 220 km/s** and  **$v_{\text{esc}} = 544 \text{ km/s}$**
- **$v_e \sim 245 \text{ km/s}$**  - WIMP velocity relative to Earth
- Local density of WIMPs =  **$0.3 \text{ GeV/cm}^3$**

*If WIMPs are  $100 \text{ GeV}/c^2$  particles, then  $\sim 10$  million pass through your hand each second!*



# Scattering rate dissected

$$\begin{array}{c}
 \text{Interaction Rate} \\
 \text{[events/keV/kg/day]}
 \end{array}
 \frac{dR}{dE_R} =
 \begin{array}{c}
 \text{particle} \\
 \text{theory}
 \end{array}
 \frac{\sigma_o}{m_\chi}
 \begin{array}{c}
 \text{nuclear} \\
 \text{structure}
 \end{array}
 \frac{F^2(E_R)}{m_\tau^2}
 \begin{array}{c}
 \text{local properties} \\
 \text{of DM halo}
 \end{array}
 \frac{\rho_o T(E_R)}{v_o \sqrt{\pi}}$$

$\uparrow$   
 Recoil energy of nucleus

# Scattering rate dissected

		particle theory	nuclear structure	local properties of DM halo
Interaction Rate [events/keV/kg/day]	$\frac{dR}{dE_R} =$	$\frac{\sigma_o}{m_\chi}$	$\frac{F^2(E_R)}{m_r^2}$	$\frac{\rho_o T(E_R)}{v_o \sqrt{\pi}}$

$$\sigma_o \simeq \frac{4m_r^2}{\pi} f A^2 \leftarrow$$

*For spin-independent scattering, and small momentum transfer, scattering terms add coherently, proportional to  $A^2$  ( $A$ = atomic mass)*

*WIMP-nucleon coupling constant;  
assumed same for proton and  
neutron in vanilla scattering*

# Scattering rate dissected

$$\begin{array}{c} \text{Interaction Rate} \\ \text{[events/keV/kg/day]} \end{array} \quad \frac{dR}{dE_R} = \overset{\text{particle theory}}{\boxed{\frac{\sigma_o}{m_\chi}}} \overset{\text{nuclear structure}}{\boxed{\frac{F^2(E_R)}{m_r^2}}} \overset{\text{local properties of DM halo}}{\boxed{\frac{\rho_o T(E_R)}{v_o \sqrt{\pi}}}$$

$$m_r = \frac{m_\chi m_N}{m_\chi + m_N} \quad \text{Reduced mass of WIMP-nucleon system}$$

Form factor parameterizes  
“coherence” vs  $E_r$

$$F(E_R) \simeq \exp \left( -E_R m_N R_o^2 / 3 \right)$$

# Scattering rate dissected

		particle theory	nuclear structure	local properties of DM halo
Interaction Rate [events/keV/kg/day]	$\frac{dR}{dE_R} =$	$\frac{\sigma_o}{m_\chi}$	$\frac{F^2(E_R)}{m_r^2}$	$\frac{\rho_o T(E_R)}{v_o \sqrt{\pi}}$

*Integral over local WIMP velocity distribution (Maxwell-boltzmann w/ assumed parameters on earlier slide)*

$$T(E_R) \simeq \int_{v_{min}}^{\infty} \frac{f(\mathbf{v}_D, \mathbf{v}_E, v_{esc})}{v_D} dv_D$$


$$v_{min} = \sqrt{E_R m_N / (2m_r^2)}$$

*$v_{min}$  is the minimum WIMP velocity needed to produce recoil  $E_r$*

# The expected signal from a heavy WIMP

*Expected recoil spectrum is roughly exponential with  $\ll 1$  event/kg/day expected,  $A^2$  enhancement helps a lot with heavy WIMPs*


100 GeV/c<sup>2</sup> WIMP-induced recoil spectrum




# The expected signal from a light WIMP

*Recoil spectrum drops off much more steeply with energy because kinematics matter much more for light WIMPs !*

10 GeV/c<sup>2</sup> WIMP-induced recoil spectrum



A WIMP must have a minimum velocity to produce a recoil of a specific energy



➔ *Experiments with lighter targets and lower thresholds have the advantage when looking for WIMPs with mass < 10 GeV/c<sup>2</sup>*



# Backgrounds: as big a problem as ever

*Expected WIMP scattering rate is  $\sim 10^7$  times lower than radioactivity of common materials*

## ELECTRON RECOILS

Gamma: MOST PREVALENT BACKGROUND

Beta: most common “surface events”



## NUCLEAR RECOILS


Neutron: rare but single-scatters NOT distinguishable from a WIMP signal

Alphas: not a background for CDMS


Pb recoils (from alpha decay): another type of surface event background

Have we seen evidence  
for WIMPs?


# Spin-Independent Landscape




# Spin-Independent Landscape



# Zoom of Low-Mass Region




# Hint from CDMS II Silicon Search



PRL 111 (2013) 251301

*Likelihood analysis incorporating energy of events yields  $\sim 3\sigma$  significance*




# Recent updates to earlier hint: CoGeNT

In 2010, CoGeNT using PPC Ge to push ionization thresholds down to  $<0.5$  keV; reported an excess of low-energy events with spectrum consistent with a  $\sim 10$  GeV/ $c^2$  WIMP


In 2011, reports a modulation of events in the 0.5-3.0 keVee region with  $\sim 2\sigma$  significance, corresponding to a large fractional modulation

In 2014, Analysis of 3.4 years of data shows persistent  $\sim 2\sigma$  modulation in low-energy region, [arXiv:1401.3295](#); Alternative maximum likelihood analysis qualitatively supports earlier analysis, but with less significant excess seen at low energies, [arXiv:1401.6234](#).



# (Ultra) Low Ionization Threshold Experiment: CDMSlite


Neganov-Luke amplification of phonon response allows operation at very low energy thresholds




Electrons and holes radiate phonons proportional to  $V_{\text{bias}}$  as they drift to the electrodes. → Apply large  $V_{\text{bias}}$  to amplify ionization signal

# (Ultra) Low Ionization Threshold Experiment: CDMSlite

Neganov-Luke amplification of phonon response allows operation at very low energy thresholds



Electrons and holes radiate phonons proportional to  $V_{\text{bias}}$  as they drift to the electrodes. → Apply large  $V_{\text{bias}}$  to amplify ionization signal





*First CDMSlite run: 170 eVee ( $<1 \text{ keV}_{\text{nr}}$ ) threshold with 0.6 kg Ge, 10 live days and no background subtraction!*



# First Results from LUX

85.3 live days with  
118 kg Xe target, operated in  
dual-phase TPC; Sets world's most  
sensitive SI limit over broad mass range



*Large enough mass  
can give sensitivity  
to some low mass  
WIMPs. But be  
careful with energy  
scale calibration and  
velocity profiles!*


# Why low mass?

*Masses  $< 10 \text{ GeV}/c^2$  are not naturally preferred by many theoretical frameworks motivated by the WIMP miracle. However ....*

- Many models predict dark matter outside of the “vanilla” WIMP paradigm. Fine tuning of parameters is often necessary, even if it’s undesirable
- Expanding beyond CMSSM (even SUSY) opens up a lot of parameter space: *pMSSM, NMSSM, Asymmetric, Isospin Violating, Inelastic, (insert your favorite model here), ...*
- We should not ignore the data. Several experiments are reporting excess events. Could these be the first indications of a major discovery? Several other experiments, done with different targets, are in tension with a dark matter interpretation...

Even if the experiments are only seeing backgrounds, its worth gathering enough data to definitively rule out these anomalous observations!


# SuperCDMS Low-Mass WIMP Search




# SuperCDMS Soudan



*interleaved*  
*Z*-sensitive  
*I*onization &  
*P*honon detector




*Improved fiducialization from measurement  
of z-symmetric ionization response  
Phonon guard and z-symmetric phonon  
response helps too!*




15 germanium detectors  
0.6 kg each  
Operational since March of 2012

**Data for this analysis:** 577 kg-days  
taken from Mar 2012 – July 2013  
7 iZIPs w/ lowest trigger thresh



APL 103, 164105(2013)



# The SuperCDMS Collaboration



California Inst. of Tech.  
B. Cornell, S.R. Golwala,  
R.H. Nelson



CNRS-LPN  
Q. Dong, A. Cavanna,  
U. Gennser, L. Couraud, Y. Jin



FNAL  
R. Basu Thakur, D.A. Bauer,  
D. Holmgren, L. Hsu, B. Loer



Mass. Inst. of Tech.  
A.J. Anderson, J. Billard,  
E. Figueroa-Feliciano,  
A. Leder, K.A. McCarthy



NIST Inst. of Tech.  
J. Ullom



PNNL  
J. Hall



Queen's University  
C.H. Crewdson, P.C.F. Di Stefano,  
O. Kamaev, P. Nadeau, K. Page,  
W. Rau, Y. Ricci



Santa Clara University  
B.A. Young



SLAC  
M. Asai, A. Borgland, D. Brandt,  
P.L. Brink, G.L. Godfrey,  
M.H. Kelsey, R. Partridge,  
K. Schneck, D.H. Wright



Southern Methodist U.  
R. Calkins, J. Cooley, B. Kara,  
H. Qiu, S. Scorza



Stanford University  
B. Cabrera, D.O. Caldwell,  
R.A. Moffatt, P. Redl,  
B. Shank, S. Yellin, J.J. Yen



Syracuse University  
M.A. Bowles, R. Bunker,  
Y. Chen, R.W. Schnee



Texas A&M University  
H.R. Harris, A. Jastram,  
R. Mahapatra, J.D. Morales Mendoza,  
K. Prasad, D. Toback, S. Upadhyayula,  
J.S. Wilson, S. Yeager



U. Autónoma de Madrid  
D.G. Cerdeno, L. Esteban,  
E. Lopez Asamar



U. British Columbia  
S.M. Oser, W.A. Page,  
H.A. Tanaka



U. California, Berkeley  
M. Daal, T. Doughty,  
N. Mirabolfathi, A. Phipps,  
M. Pyle, B. Sadoulet, B. Serfass,  
D. Speller



U. Colorado Denver  
M.E. Huber



U. Evansville  
A. Reisetter



U. Florida  
R. Agnese, D. Balakishiyeva,  
T. Saab, B. Welliver



U. Minnesota  
D. Barker, H. Chagani, P. Cushman,  
S. Fallows, T. Hofer, A. Kennedy,  
K. Koch, V. Mandic, M. Pepin,  
H. Rogers, A.N. Villano, J. Zhang




U. South Dakota  
J. Sander



# Optimizing for Low Mass

*Recall: experiments with lighter targets and lower thresholds have the advantage when looking for WIMPs with mass  $< 10 \text{ GeV}/c^2$*



## Our strategy:

Ge is a relatively heavy target so go as low in threshold as possible



→ trigger threshold (1.6 keV $_{\text{nr}}$ )

Backgrounds more difficult to reject below 10 keV $_{\text{nr}}$ ; use full capability of iZIPs to reject as much background as possible

*We expect background events in the signal region!! Tradeoff is greater sensitivity to low mass WIMPs.*

# Backgrounds to Eliminate



**Bulk electron recoils =**  
Compton background and 1.3  
keV activation line



*Ionization vs phonon  
distinguishes NR  
from bulk ER*


# Backgrounds to Eliminate

**Bulk electron recoils =**  
Compton background and 1.3  
keV activation line



*Ionization vs phonon  
distinguishes NR  
from bulk ER*

**sidewall & surface events =**  
betas and x-rays from  $^{210}\text{Pb}$ ,  $^{210}\text{Bi}$ ,  
recoils from  $^{206}\text{Pb}$  (i.e. Rn  
daughters), outer radial compton  
and ejected electrons from  
compton scattering





*Use division of energy  
between inner and outer  
sensors, "radial partition"*



*Use division of energy  
between sides 1 and 2,  
"z-partition"*



# Backgrounds to Eliminate

**Bulk electron recoils =**  
Compton background and 1.3  
keV activation line



*Ionization vs phonon  
distinguishes NR  
from bulk ER*

**sidewall & surface events =**  
betas and x-rays from  $^{210}\text{Pb}$ ,  $^{210}\text{Bi}$ ,  
recoils from  $^{206}\text{Pb}$  (i.e. Rn  
daughters), outer radial comptons  
and ejected electrons from  
compton scattering



*Use division of energy  
between inner and outer  
sensors, “radial partition”*

*Use division of energy  
between sides 1 and 2,  
“z-partition”*


**Cosmogenic & radiogenic  
neutrons**



*Use active and passive  
shielding. Simulation  
determines remaining  
irreducible rate*

# Nuclear Recoil Energy Determination

*Ionization for nuclear recoils,  
measured from  $^{252}\text{Cf}$  data:*



*Total phonon energy =*

$$E_{\text{total}} = E_{\text{luke}} + E_{\text{recoil}}$$

*$E_{\text{total}}$  is measured with phonons*

*NR equivalent energy =*

$$E_{\text{total}} - E_{\text{Luke NR}}$$

*$E_{\text{Luke NR}}$  estimated from mean NR  
ionization, varies with  $E_{\text{total}}$   
(same as CDMS II low mass  
search)*


*Note: we sometimes approximate mean ionization with Lindhard theory because measured values are detector-dependent. This is labeled “**Lindhard nuclear recoil energy**”; difference is a few %.*

# Analysis Summary

**Blind analysis:** All singles in analysis energy range removed from study, except data following neutron calibration due to activation (additional 97 kg-days not used for limit calculation or cut tuning)

# Analysis Summary

**Blind analysis:** All singles in analysis energy range removed from study, except data following neutron calibration due to activation (additional 97 kg-days not used for limit calculation or cut tuning)



**Efficiencies:** measured with neutrons from  $^{252}\text{Cf}$ . Geant4 used to correct for multiple scattering, yields ~25% correction

## Data Quality:

- Reject high/abnormal noise
- Reject atypical operational periods

## Trigger and Analysis Threshold:

- Select periods of stable, well-defined trigger threshold
- Analysis thresholds based on time-varying noise baseline

## Preselection:


- Single-detector scatter
- Muon veto anticoincident
- ionization fiducial volume
- Ionization energy and phonon partitions consistent w/ NR

## Boosted Decision Tree

- “tight” phonon fiducial volume and ionization yield at low energy

# Boosted Decision Tree (BDT)


*Discrimination lies in correlations between 4 parameters in partition and energy*



WIMP model assumes:


$$\sigma = 6 \times 10^{-42} \text{ cm}^2$$

# Background model w/ pulse simulation



***Problem:*** Backgrounds at low energy are more difficult to separate from signal region due to worsening resolution


# Background model w/ pulse simulation



**Problem:** Backgrounds at low energy are more difficult to separate from signal region due to worsening resolution

**Solution:** Study directly with a pulse simulation; using high energy events in sidebands and calibration data as templates

Event w/ good signal to noise, scaled down in amplitude




Random trigger (e.g. noise)



+

=

Simulated low energy event



reconstruction software



weight events as a function of energy to match low energy backgrounds

# Background estimates

Background estimates finalized before unblinding, included known systematic effects; Checked against open dataset and reasonable agreement found

\*Purpose of background model was tuning cuts; possible unknown systematics preclude background subtraction for this blind analysis.

*Thus, decision made to set an upper limit prior to unblinding*


*4 BDT cuts developed for 5, 7, 10 and 15 GeV/c<sup>2</sup> WIMPs; accept events that pass any of the four cuts; Each cut was tuned simultaneously on all detectors, maximizing 90% C.L. poisson sensitivity for that mass*

**Background model expected:  $6.1^{+1.1}_{-0.8}$  events**

**Neutron background adds additional:  $0.10 \pm 0.02$  events**


# Unblinding: *Before BDT cut*

*Shown: events passing all cuts except the BDT and ionization selection*




# Unblinding: *After BDT cut*

*11 candidates seen,  $6.2 +1.1 -0.8$  expected*







# w/ 95% WIMP Confidence Intervals

*11 candidates seen,  $6.2 +1.1 -0.8$  expected*



# How model compares to data post-unblinding



WIMP model assumes:


$$\sigma = 6 \times 10^{-42} \text{ cm}^2$$

Fermilab W&C, March 2014

# Post-unblinding discussion


Events generally high in quality  
except lowest energy candidate, which looks like spurious noise.

*Agreement  
with  
predicted  
background  
is good on  
most  
detectors*




# Post-unblinding discussion

Events generally high in quality  
except lowest energy candidate, which looks like spurious noise.




Probability for background to fluctuate up to 3 or more events is 0.0004 on T5Z3. This detector has a shorted ionization guard; at present it is unclear whether excess events are related, additional studies are ongoing

# Post-unblinding discussion



# Spin-independent Scattering Constraints


*90% C.L. optimal interval upper limit, no background subtraction, treating all observed (eleven) events as WIMP candidates*



- CoGeNT strongly disfavored in model-independent scenario
- CDMS II (Si) disfavored under assumption of standard halo model and  $A^2$  coupling
- Explores new parameter space below 6 GeV/c<sup>2</sup>
- Competitive constraint for Ge up to 20 GeV/c<sup>2</sup>; dedicated HT analysis yet to come
- Disagreement between limit and sensitivity at high WIMP mass due to events on T5Z3.


# What about the CDMS II Si result?

*Available parameter space is being tightly constrained by this result and others, but some select models still remain....*






Example above from recent paper on isospin-violating dark matter  
(w/ colored mediators)

# The longer term picture (SuperCDMS G2)



# WIMP Searches w/ SuperCDMS SNOLAB

~100 kg of mixed Ge/Si payload,  
w/ 5% detectors configured in  
CDMSlite mode



- Locate in North America's deepest underground lab
- Bigger iZIP detectors
- Cleaner shielding, w/ active neutron veto
- Upgraded electronics
- Room to expand to 400 kg

# Summary

First result using background rejection capability of  
SuperCDMS!

arXiv: 1402.7137

7 iZIPs analyzed down to trigger threshold ( $1.6 \text{ keV}_{\text{nr}}$ );  
Exposure of 577 kg-days sets 90% C.L. upper limit to  
WIMP-nucleon SI scattering,  $\sigma = 1.2 \times 10^{-42} \text{ cm}^2$ , at  $8 \text{ GeV}/c^2$

*New phase space explored below  $6 \text{ GeV}/c^2$*

CoGeNT interpretation of WIMPs strongly disfavored in model-independent scenario; CDMS II (Si) region disfavored under standard halo model and  $A^2$  coupling

SuperCDMS SNOLAB will have unprecedented reach in searches for low-mass WIMPs and complementary sensitivity in searches for high-mass WIMPs